

## 23. Urine sodium

Ward	Not stated	D.O.B/Age	10/06/1958
Consultant			

Urine sodium 224 mmol/L

Request form: hyponatraemia

Many patients present due to manifestations of other medical comorbidities, with hyponatremia being recognized only secondarily. Many medical illnesses, such as chronic heart failure, liver failure, renal failure, or pneumonia, may be associated with hyponatremia. Patients usually present with symptoms related to their primary illness.

Symptoms of hyponatremia range from nausea and malaise, with a mild reduction in the serum sodium, to lethargy, decreased level of consciousness, headache, and (if severe) seizures and coma. Overt neurologic symptoms most often are due to very low serum sodium levels (usually < 115 mmol/L), resulting in intracerebral osmotic fluid shifts and brain oedema.

Examination should include orthostatic vital signs and an accurate assessment of volume status. Volume status forms an integral part of assessment as it often guides assessment and treatment.

A full assessment for medical comorbidities is also essential, with particular attention to cardiopulmonary and neurologic components of the examination.

Authorised by Dr TA Gcingca on 05/01/2020 at 21:37  
Urine sodium 224 mmol/L

Authorised by Dr TA Gcingca on 05/01/2020 at 21:37  
Urine osmolality 649 mmol/kg 50 - 1200

CT brain and CXR may be indicated if SIADH suspected.

True hyponatraemia.

Hyponatremia can be classified according to volume status, as follows:

- Hypovolemic hyponatremia: decrease in total body water with greater decrease in total body sodium
- Euvolemic hyponatremia: normal body sodium with increase in total body water
- Hypervolemic hyponatremia: increase in total body sodium with greater increase in total body water

Hyponatremia can be further subclassified according to effective osmolality, as follows:

- Hypotonic hyponatremia
- Isotonic hyponatremia
- Hypertonic hyponatremia

There are three essential laboratory tests in the evaluation of patients with hyponatremia that, together with the history and the physical examination, help to establish the primary underlying etiologic mechanism: urine osmolality, serum osmolality, and urinary sodium concentration.

1. Urine osmolality: essential to differentiate a deficiency in excreting free water vs primary polydipsia. Urine osmolality greater than 100 mOsm/kg

indicates impaired ability of the kidneys to dilute the urine.

2. Serum osmolality: differentiates between true hyponatremia and pseudohyponatremia. True hyponatraemia causes an decrease in serum osmolality.
3. Urinary sodium: helps to differentiate between hyponatremia secondary to hypovolemia and the syndrome of inappropriate ADH secretion (SIADH). With SIADH (and salt-wasting syndrome), the urine sodium is greater than 20-40 mmol/L. With hypovolemia, the urine sodium typically measures less than 25 mmol/L.

Ancillary testing may also help with differentiating SIADH from salt-wasting. Serum uric acid levels can be important supportive information (they are typically reduced in SIADH and also reduced in salt wasting). After correction of hyponatremia, the hypouricemia corrects in SIADH but remains with a salt-wasting process.

---

## 7. EDTA contamination vs renal impairment

Ward	Surgical ICU	D.O.B/Age	17/04/1994
Consultant			

Potassium: 6.1 H mmol/L [3.5 – 5.1]

No diagnosis on request form, unable to get hold of clinician.

Authorised by Dr TA Gcingca on 27/11/2019 at 08:37

Sodium	137	mmol/L	136 - 145
--------	-----	--------	-----------

Authorised by Dr TA Gcingca on 27/11/2019 at 08:37

Potassium	6.1 H	mmol/L	3.5 - 5.1
-----------	-------	--------	-----------

Authorised by Instrument on 27/11/2019 at 06:11

Chloride	106	mmol/L	98 - 107
----------	-----	--------	----------

Authorised by Dr TA Gcingca on 27/11/2019 at 08:37

Urea	19.7 H	mmol/L	2.1 - 7.1
------	--------	--------	-----------

---

Authorised by Instrument on 27/11/2019 at 06:11

Creatinine	198 H	umol/L	64 - 104
eGFR (MDRD formula)	38	mL/min/1.73 m <sup>2</sup>	

MDRD-derived estimation of GFR may significantly underestimate true GFR in patients with GFR > 60 mL/min/1.73m<sup>2</sup>. It may also be unreliable in the case of: age <18 years or >70 years; pregnancy; serious co-morbid conditions; acute renal failure; extremes of body habitus/unusual diet; gross oedema. The MDRD-eGFR used here does not employ an ethnic factor for race.

Authorised by Dr TA Gcingca on 27/11/2019 at 08:37

Calcium	1.17 L	mmol/L	2.15 - 2.50
---------	--------	--------	-------------

Authorised by Dr TA Gcingca on 27/11/2019 at 08:37

Magnesium	0.97	mmol/L	0.63 - 1.05
-----------	------	--------	-------------

Authorised by Instrument on 27/11/2019 at 06:11

Inorganic phosphate	1.46 H	mmol/L	0.78 - 1.42
---------------------	--------	--------	-------------

Authorised by Instrument on 27/11/2019 at 06:11

Indices in serum:

Haemoglobin index	Not detected
Bilirubin index	Trace
Lipaemia index	Not detected

Authorised by Instrument on 27/11/2019 at 05:44

White Cell Count	10.17	x 10 <sup>9</sup> /L	3.92 - 10.40
Red Cell Count	3.32 L	x 10 <sup>12</sup> /L	4.50 - 5.50
Haemoglobin	9.8 L	g/dL	13.0 - 17.0
Haematocrit	0.274 L	L/L	0.400 - 0.500
MCV	82.5 L	fL	83.1 - 101.6
MCH	29.5	pg	27.8 - 34.8
MCHC	35.8 H	g/dL	33.0 - 35.0
Red Cell Distribution Width	15.2	%	12.1 - 16.3
Platelet Count	116 L	x 10 <sup>9</sup> /L	171 - 388

Potassium ethylenediaminetetraacetic acid (EDTA) is a sample tube anticoagulant used for many laboratory analyses. Gross potassium EDTA contamination of blood samples is easily recognised by marked hyperkalaemia and hypocalcaemia. Subtle contamination is a relatively common, often unrecognised erroneous cause of spurious hyperkalaemia. In the case illustrated, it would be difficult to confidently exclude EDTA contamination based on these results alone. There is renal impairment which may explain the hyperkalaemia. The increased phosphate coupled with the renal impairment would also be an argument for the hypocalcaemia present.

In this instance, comparison with previous results was useful. The results are most likely due to renal impairment. As the patient had been admitted to the ward for a week, it was useful to be able to compare previous results. The gradual decline in renal function helped to explain the biochemical findings. As the samples were drawn on different days by different persons, the likelihood of EDTA contamination on all the days is relatively slim.

However, it is important to be cognisant that mild EDTA contamination may cause subtle shifts in results that may have negative consequences for the patient if erroneously acted on.

---

## 24. CoA trapping

<b>Ward</b>	Paeditric ICU	<b>D.O.B/Age</b>	11/03/2020
<b>Consultant</b>	Prof G. vd Watt		

Elevated propionic acid in the urine organic acid profile.

Fever with LRTI. ?COVID

Normal birth with no antenatal problems

#RVD exposed

Now:

#FTT

#LRTI. ?COVID

The patient presented with fever and LRTI which resolved after 3 -4 days of antibiotics. The patient then developed seizures with apnoeic attacks. The patient required intubation and ventilation and was transferred to ICU. The patient was noted to be having breakthrough seizures despite anticonvulsant therapy.

Further questioning revealed that the patient had become progressively drowsy with poor feeding.

Family history: No siblings noted to have had previous problem.

The patient was noted as not interacting with his environment.

CNS exam: Low GCS with upper motor neuron signs.

Other systems unremarkable.

pH	7.13 L		7.35 - 7.45
pCO2	2.99 L	kPa	4.66 - 6.38
pO2	19.90 H	kPa	11.04 - 14.36
Standard bicarbonate	9 L	mmol/L	22 - 26
Base excess	-21.6 L	mmol/L	-10.0 - -2.0
O2 saturation	100 H	%	94 - 98
Sodium	121 L	mmol/L	136 - 145
Potassium	4.4	mmol/L	3.5 - 4.5
Chloride	92 L	mmol/L	98 - 113
Glucose	13.3	mmol/L	
Ionised calcium	0.80	mmol/L	
Carboxyhaemoglobin	3.5	%	
Methaemoglobin	-1.7	%	

Authorised by NL Makhalima on 28/05/2020 at 16:42

Ammonia	1517 H	umol/L	40 - 80
---------	--------	--------	---------

Please note that preanalytical factors including a delay in sample reception and sample not transported on ice may cause raised ammonia results.

Trace lipaemia observed

Please repeat

Total cholesterol	1.90	mmol/L
Triglyceride	6.21	mmol/L
HDL cholesterol	0.18	mmol/L
LDL cholesterol	Triglyceride level too high [>4.5mmol/l] for LDL calculation	

CHOLESTEROL TREATMENT TARGETS (per CV Event Risk Category):

Risk Category:	TC target:	LDL-C target:
Low/Moderate Risk	<5.0 mmol/L	<3.0 mmol/L
High Risk	<4.5 mmol/L	<2.5 mmol/L

Authorised by KF Sephula on 28/05/2020 at 05:29

Calcium	1.24 L	mmol/L	2.12 - 2.64
---------	--------	--------	-------------

Authorised by KF Sephula on 28/05/2020 at 05:29

Albumin	23 L	g/L	26 - 41
---------	------	-----	---------



Authorised by NL Makhalima on 29/05/2020 at 14:07			
Total bilirubin	5	umol/L	5 - 21
Authorised by NL Makhalima on 29/05/2020 at 14:07			
Conjugated bilirubin (DBil)	2	umol/L	0 - 5
Authorised by NL Makhalima on 29/05/2020 at 17:52			
Alanine transaminase (ALT)	106 H	U/L	1 - 25
Authorised by NL Makhalima on 29/05/2020 at 14:08			
Aspartate transaminase (AST)	391 H	U/L	0 - 51
Authorised by NL Makhalima on 29/05/2020 at 14:08			
Alkaline phosphatase (ALP)	382 H	U/L	75 - 316
Authorised by NL Makhalima on 29/05/2020 at 14:07			
Gamma-glutamyl transferase (GGT)	44	U/L	12 - 122

Authorised by B Gool on 26/05/2020 at 16:35

CSF glucose 1.5 mmol/L

CSF glucose reference range:

CSF glucose is normally 60 - 80% of plasma glucose, in samples taken within 15 minutes of each other.

Authorised by B Gool on 26/05/2020 at 16:35

CSF protein 1.62 H g/L 0.20 - 0.80

Authorised by NL Makhalima on 26/05/2020 at 17:50

CSF adenosine deaminase 0.0 U/L

CSF ADA activity of > 6 U/L is suggestive of TB. However, other conditions such as bacterial or Cryptococcal meningitis may also produce elevated ADA levels.

**CSF Analysis:****Appearance:**

Clarity	Bloodstained
Clots	Absent

**Cell Count:**

Polymorphs	0	/uL
Lymphocytes	0	/uL
Erythrocytes	48	/uL

Authorised by NT Jikwana on 26/05/2020 at 14:53

**Gram Stain:**

Organisms	No bacteria observed
-----------	----------------------

Authorised by MG Mpotje on 28/05/2020 at 09:07

**Bacterial Culture:**

No growth after 2 days

---

Authorised by NL Makhalima on 28/05/2020 at 16:45

White Cell Count	0.59 L	x 10 <sup>9</sup> /L	5.00 - 20.00
Red Cell Count	2.54 L	x 10 <sup>12</sup> /L	3.90 - 5.90
Haemoglobin	8.1 L	g/dL	12.0 - 21.8
Haematocrit	0.218 L	L/L	0.340 - 0.620
MCV	85.7 L	fL	88.0 - 126.0
MCH	31.7	pg	31.0 - 37.0
MCHC	37.0 H	g/dL	30.0 - 36.6
Red Cell Distribution Width	14.8	%	
Platelet Count	67 L	x 10 <sup>9</sup> /L	140 - 350
MPV	9.6	fL	7.0 - 11.4

Comment	Automated platelet count to be reviewed microscopically.
---------	---

MCHC results may be affected by lipaemia

repeated tplateet = 71

FBC comment:

No clot detected in EDTA sample

Peripheral smear to be reviewed

CT brain may be useful in assess for organic neurological

cause.

Propionic acidaemia.

DDx: Biotinidase deficiency

Propionic acidaemia is an organic acidaemia characterized by deficiency of propionyl-CoA carboxylase. Propionyl-CoA carboxylase converts propionyl-CoA to methylmalonyl-CoA. It is inherited in an autosomal recessive pattern. The metabolism of isoleucine, valine, threonine, and methionine produces propionyl-CoA. To a lesser degree, cholesterol and odd-chain fatty acids also contribute to propionyl-CoA levels. Affected individuals must follow a low-protein diet and early diagnosis improved prognosis.

The accumulation of propionyl-CoA results in significant mitochondrial CoA trapping and inhibited fatty acid oxidation. The enhanced anapleurosis of propionate and CoA trapping alters the pool sizes of tricarboxylic acid cycle (TCA) metabolites. This explains the marked hyperammonaemia that patients present with as well as potential hypoglycaemia

A high index of suspicion is required to diagnose inborn errors of metabolism (IEM). This case highlighted the importance of understanding key points in metabolic pathways. It also emphasized the correlation between catabolic stress being an initiating event in IEMs.

---

## 22. Haemolysis (intravascular)

Ward	Surgical ward	D.O.B/Age	26 y.o
------	---------------	-----------	--------

<b>Consultant</b>	Dr H. Vreede		
-------------------	--------------	--	--

#### Indices in serum:

Haemoglobin index	4+
Bilirubin index	Trace
Lipaemia index	Trace

Call from a clinician to assist with generating results that were being rejected due to haemolysis.

26 y.o male

#Previously well

Admitted with multiple stab wounds and had a haemo-pneumothorax on the left side. Intercostal chest drain inserted. The patient acutely decompensated after 3 days being admitted. He was noted to have metabolic acidosis, hyperlactataemia, and symptoms of shock. The patient was not on any medication besides analgesia. No previous blood transfusion. No procedure in the ward.

Patient noted to be jaundiced. Urine coke-coloured. No petechiae. Patient not bleeding from any wound sites.

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Sodium	127 L	mmol/L	136 - 145
--------	-------	--------	-----------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Potassium	6.5 H	mmol/L	3.5 - 5.1
-----------	-------	--------	-----------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Urea	20.0 H	mmol/L	2.1 - 7.1
------	--------	--------	-----------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Creatinine	272 H	umol/L	64 - 104
eGFR (MDRD formula)	25	mL/min/1.73 m <sup>2</sup>	

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Calcium	2.15	mmol/L	2.15 - 2.50
---------	------	--------	-------------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Magnesium	0.87	mmol/L	0.63 - 1.05
-----------	------	--------	-------------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Inorganic phosphate	2.05 H	mmol/L	0.78 - 1.42
---------------------	--------	--------	-------------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Albumin	24 L	g/L	35 - 52
---------	------	-----	---------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Total bilirubin	44 H	umol/L	5 - 21
-----------------	------	--------	--------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Conjugated bilirubin (DBil)	3	umol/L	0 - 3
-----------------------------	---	--------	-------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Alanine transaminase (ALT)	342 H	U/L	10 - 40
----------------------------	-------	-----	---------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Alkaline phosphatase (ALP)	88	U/L	53 - 128
----------------------------	----	-----	----------

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Haptoglobin	0.46	g/L	0.30 - 2.00
-------------	------	-----	-------------

Authorised by Instrument on 06/11/2019 at 04:38

Indices in serum:


Haemoglobin index	4+
Bilirubin index	Trace
Lipaemia index	Trace

Authorised by Dr TA Gcingca on 06/11/2019 at 14:21

Chemistry comment:

Gross haemolysis is present. The decreased haptoglobin level points toward intravascular haemolysis but please treat results with reserve and correlate with clinical findings.





Patient sample after centrifugation



Gross haemolysis

CXR: haemopneumothorax on the left.

?sepsis ?toxin introduced through stab wounds

Intravascular haemolysis results in the release of cell-free haemoglobin, red blood cell (RBC) stroma, and non-stroma proteins. Free haemoglobin binds nitric oxide (NO) at rate 1000 times that of RBC. Haemoglobin scavenging leads to decreased bioavailability of NO and thus vasoconstriction and alterations in capillary response to hypoxia. RBC stroma, which is the cytoskeletal framework supporting haemoglobin, can also contribute to DIC pathogenesis via activation of platelets and coagulation cascade. RBC stroma has also been shown to increase blood pressure and is toxic to the glomerulus and renal tubule and thus can cause acute renal failure. Ultimately, increased cytokines and hypotension stimulate a compensatory sympathetic nervous system response contributing to renal, splanchnic, and cutaneous

vasoconstriction that, in combination with pathophysiology described above, leads to shock and circulatory collapse.

Marked increase of lactate dehydrogenase and haemosiderinuria are typical of intravascular haemolysis. Several haemolytic markers are available to guide the differential diagnosis and to monitor treatment of haemolytic conditions. They include increased reticulocytes (an indicator of marrow compensatory response) elevated lactate dehydrogenase, reduced haptoglobin, and unconjugated hyperbilirubinemia.

However, increased reticulocytes, lactate dehydrogenase, and bilirubin, as well as reduced haptoglobin, are observed in conditions other than haemolysis that may confound the clinical picture.

Haptoglobin is a positive acute-phase reactant. It is a protein that binds irreversibly to free (oxy)haemoglobin liberated into the plasma during intravascular haemolysis. The haptoglobin-haemoglobin complex is removed rapidly by the reticuloendothelial system to prevent loss of haemoglobin in urine. Low levels are a diagnostic indicator of intravascular haemolysis (but may be low in liver disease or with endogenous or exogenous oestrogen). Elevated levels are associated with acute phase response, nephrotic syndrome and with corticosteroids.

It is interesting to note in this patient that his result is in the lower level of normal, pointing towards the possibility that haptoglobin may be markedly decreased

---

## 4. PSA

Ward	Casualty Department	D.O.B/Age	04/12/1940
Consultant			

Prostate-specific Ag (PSA) **949.50 H**  
ug/L **<4.00**

Urinary retention.

Request form: Lower urinary tract symptoms and urinary retention.

Important clinical findings to assess for include:

**General:** Temporal wasting, signs of urinary incontinence (e.g. any leaking noted, need to wear sanitary products)

**Abdominal:** Assess for masses, palpable bladder from retention

**P.R:** Assess prostate for size, consistency, tenderness.

**CNS:** Assess for any neurological fallout as prostatic metastasis tend to metastasize to the lower vertebrae.

Creatinine **83**  
umol/L **64 – 104**  
eGFR (MDRD formula) **>60**  
mL/min/1.73 m<sup>2</sup>

White Cell Count **5.01** **x**  
10<sup>9</sup>/L **3.92 – 10.40**  
Red Cell Count **5.39** **x**  
10<sup>12</sup>/L **4.50 – 5.50**  
Haemoglobin **15.5**  
g/dL **13.0 – 17.0**  
Haematocrit **0.485**  
L/L **0.400 – 0.500**  
MCV **90.0**  
fL **83.1 – 101.6**  
MCH **28.8**  
pg **27.8 – 34.8**

MCHC		32.0	L
g/dL	33.0 – 35.0		
Red Cell Distribution Width		13.2	
%	12.1 – 16.3		
Platelet Count		226	x
10 <sup>9</sup> /L	171 – 388		

#### PATHOLOGICAL DIAGNOSIS:

Prostate, biopsy: Adenocarcinoma.

Imaging studies may be necessary if there is a concern for metastasis and these will be guided by the clinical presentation e.g. CXR if metastasis to the lungs is suspected vs MRI if there is a concern of vertebral collapse.

Prostatic adenocarcinoma.

- Prostate-specific antigen (PSA) is a protein produced by normal prostatic cells. The majority of PSA is produced by the glands in the transitional zone of the prostate (BPH). The peripheral zone, where 80% of prostate cancers originate, produces very little PSA.
- An enlarged prostate can cause obstructive uropathy. The creatinine values in this patient do not suggest renal impairment though a baseline creatinine would be required to assess this.
- PSA is used for screening, diagnosis as well as monitoring of prostate related disease processes. PSA is an organ-specific, not a cancer-specific marker. It is useful in detection, staging and monitoring of prostate cancer.
- To improve diagnostic accuracy when PSA is between 4-10ug/L ("grey zone"), free PSA is measured and the free/total PSA ratio is calculated. Most normal PSA is protein-bound, and in prostatic cancer, a greater proportion is unbound. A free/total PSA ratio <0.25 increases the likelihood of cancer.

---

## 21. APT

<b>Ward</b>	Maternity Ward	<b>D.O.B/Age</b>	27 y.o
<b>Consultant</b>	Dr C. Hudson		

APT test positive.

?haemolytic disease of the newborn

27 y.o. female

G2P1 at 34 weeks

RH negative with Rhesus iso-immunization. Anti-D titres 1:128. (Blood group AB negative). Coombs test positive. Risk of haemolytic disease of the new-born.

Not applicable/difficult to examine foetus in-utero. Ultrasound of the middle cerebral artery peak systolic velocity was suggestive that the baby was anaemic.

U/s guided chordocentesis done and foetal blood sample obtained in utero. FBC whilst in utero showed Hb: 9.8.

Clinician requested APT test to ensure that foetal sample obtained during chordocentesis.

Haematology did Kleihauer Betke Test and it showed 100% foetal haemoglobin. APT test also correlated and showed foetal haemoglobin.

Ultrasound: middle cerebral artery peak systolic velocity suggestive that the baby was anaemic. U/s guided chordocentesis done and foetal blood sample obtained in utero. FBC whilst in utero showed Hb: 9.8.

APT test confirmed that foetal blood had been obtained during

chordocentesis. It also correlated with the Kleihauer Betke done by haematology.




Image 1

HbA: adult haemoglobin; HbF: foetal haemoglobin; PtC: patient control; PtT: test control.

Note the slight green tinge of HbA

Principles of the APT test: Sodium hydroxide (NaOH) denatures adult oxyhaemoglobin to haematin (with a colour change from pink to yellow green). Foetal haemoglobin resists alkaline denaturation by NaOH and maintains a pink colour. If adult haemoglobin (HbA) is present in the sample, it turns yellow and then green within two minutes of the addition of sodium hydroxide. Any pink colour that persists for longer than 2 minutes indicates foetal haemoglobin (HbF) is present in the sample.

The Kleihauer Betke Test is an acid-elution assay performed on maternal blood to determine the amount of HbF that has passed into maternal circulation. The process exposes maternal blood smear to an acid solution. HbF, being resistant to the acid, remains intact, whereas HbA is removed. Following this, the smear is stained via Shepard's method. The foetal red blood cells are left rose-pink in colour, and the maternal cells appear "ghost-like" due to the absence of staining. This test is done by Haematology. Other ways of differentiating maternal from foetal blood is examining the MCV. The foetus has a larger MCV. This is nonspecific though as the mother may have macrocytic anaemia.

---

## 3. Hyperammonaemia

<b>Ward</b>	Medical ward	<b>D.O.B/Age</b>	16/07/1950
<b>Consultant</b>			

Ammonia

umol/L

[11 – 35]

251 H



Specimen request form has hepatic encephalopathy written as the diagnosis/reason for request.

Unable to obtain.

Unable to obtain.

Sodium	143	mmol/L	[136 – 145]
Potassium	4.4	mmol/L	[3.5 – 5.1]
Urea	<b>28.1 H</b>	mmol/L	[2.1 – 7.1]
Creatinine	<b>359 H</b>	umol/L	[64 – 104]
eGFR (MDRD formula)	15	mL/min/1.73 m2	
INR		2.77	
Total bilirubin	<b>54 H</b>	umol/L	[5 – 21]
Conjugated bilirubin (DBil)		<b>36 H</b>	umol/L
		[0 – 3]	
Alanine transaminase (ALT)		34	U/L
		[10 – 40]	
Aspartate transaminase (AST)		<b>113 H</b>	U/L [15 – 40]
Alkaline phosphatase (ALP)		105	U/L
		[53 – 128]	
Gamma-glutamyl transferase (GGT)		33	
		U/L <68	
Unremarkable CMP. Elevated WCC of 11 otherwise normal FBC. Further investigations not requested on Trakcare. ?patient discharge vs transfer vs demise			

?Fulminant liver cirrhosis

?End stage liver disease

Most ammonia dealt with by the liver is produced by gut organisms. Protein degradation forms a smaller contribution. Ammonia in high concentrations is neurotoxic. It is detoxified by the liver to urea via means of the urea cycle, and urea is subsequently excreted in the urine. Pre-analytical factors including a delay in sample reception and sample not transported on ice may cause raised ammonia results. Other pre-

analytical factors to consider include:

- No smoking by the person collecting the sample or the patient the sample is being collected from.
- Tourniquet should not be applied tightly or for too long (no tourniquet application ideal).
- Collected in an EDTA container.
- Must reach the lab within 15 to 20 minutes of being collected on ice.
- Patient should be fasted.

This patient has mildly deranged liver function tests and a prolonged INR suggesting liver disease which may be contributing to the hyperammonaemia. The unremarkable elevation in the liver enzymes may be due to a decrease of viable hepatocytes.

---

## 20. Alpha-foetoprotein

Ward	Emergency unit	D.O.B/Age	07/08/1968
Consultant	Dr C. Hudson		

Request form: No clinical information provided

Unavailable.

Unavailable.

Sodium		130 L
mmol/L	136 – 145	
Potassium		3.7
mmol/L	3.5 – 5.1	
Urea		2.9
mmol/L	2.1 – 7.1	

Creatinine		64
umol/L	49 – 90	
eGFR (MDRD formula)	>60	mL/min/1.73
m2		

**Glycated haemoglobin (HbA1c):**

Glycated haemoglobin (NGSP)	6.5	%
Glycated haemoglobin (IFCC)	48	
mmol/mol		
Estimated average glucose (eAG)	7.8	mmol/L

Calcium	2.20
mmol/L	2.15 – 2.50

Total protein	86	H
g/L	60 – 78	

Albumin	28	L
g/L	35 – 52	

Total bilirubin	26	H
umol/L	5 – 21	

Conjugated bilirubin (DBil)	25	H
umol/L	0 – 3	

Alanine transaminase (ALT)	65	H
U/L	7 – 35	

Aspartate transaminase (AST)	444	H
U/L	13 – 35	

Alkaline phosphatase (ALP)	568	H
U/L	42 – 98	

Gamma-glutamyl transferase (GGT)	662	H
U/L	<40	

Lipase	91	H
U/L	13 – 60	

Alpha-feto protein (AFP)	5450	10.0	H
ug/L	0.0 – 7.0		

Thyroid stimulating hormone	6.61	H
-----------------------------	------	---

mIU/L	0.27 – 4.20		
Thyroxine (free T4)		13.7	
pmol/L	12.0 – 22.0		
White Cell Count		8.93	x
10 <sup>9</sup> /L	3.90 – 12.60		
Red Cell Count		2.92 L	x
10 <sup>12</sup> /L	3.80 – 4.80		
Haemoglobin		9.4 L	
g/dL	12.0 – 15.0		
Haematocrit		0.278 L	
L/L	0.360 – 0.460		
MCV		95.2	
fL	78.9 – 98.5		
MCH		32.2	
pg	26.1 – 33.5		
MCHC		33.8	
g/dL	32.7 – 34.9		
Red Cell Distribution Width		19.5 H	
%	12.4 – 17.3		
Platelet Count		246	x
10 <sup>9</sup> /L	186 – 454		

Medical Validation : (Authorise By Episode)

Option Mode Inquire Function Audit Print Help Result

Episode No. [redacted] MRN [redacted] HPRN [redacted] Stat [redacted]

[redacted] F 51 y 07/08/1968

Hos Groote Schuur Hospital wc GSH 021 404 9111 Collection 30/12/2019 13:04

Wrd C15 Emergency Unit 404 5208 / 09 Received 30/12/2019 13:18

Doc [redacted] Registered 30/12/2019 13:20

[redacted] ePR Detail

Visit Test Set(s)

[A] TREQ <M entry>  
[A] SPARE1 <A entry>  
[A] HBA1C <A entry>  
[A] CA <A entry>  
[A] LIPASE <A entry>  
[E] AFP # <M entry>(VQ:CR)  
[A] TSH <A entry>

[\* Curr.] (# In List)

Update Authorise  
Amend Fully Authorise  
Clear Cancel  
Notes Graph

Test Set	Staff Notes	Test Item	Result	Units	Normal Values	Previous Result 1	Previous Result 2	Previous Result 3	Previous Result 4	Previous Result 5
ALB		Albumin	28	g/L	35 - 52					
TBIL		Total bilirubin	26	umol/L	5 - 21					
		Total bilirubin auto com								
CBIL		Conjugated bilirubin (DI	25	umol/L	0 - 3					
ALT		Alanine transaminase (U	65	U/L	7 - 35					
AST		Aspartate transaminase	444	U/L	13 - 35					
ALP		Alkaline phosphatase (U	568	U/L	42 - 98					
GGT		Gamma-glutamyl transfe	662	U/L	<40					
LIPASE		Lipase	91	U/L	13 - 60					
AFP	<input checked="" type="checkbox"/>	Alpha-feto protein (AFP	545,010.0	ug/L	0.0 - 7.0					
		Machine (Serum)	COB							
		AFP auto comment	AFPCOB							
TSH		Thyroid stimulating hor	6.61	mIU/L	0.27 - 4.20					
FT4		Thyroxine (free T4)	13.7	pmol/L	12.0 - 22.0					
SIND		Serum haemoglobin inc	0							
		Serum bilirubin index	1							
		Serum lipaemia index	0							
		Serum haemoglobin va	0.00							

Staff Notes : C136

Help

[30/12/2019 22:34] [redacted] AFP Checked.

R1 - >1210  
R2 - >60500 1:50  
R3 - >121000 1:100  
R4 - >484000 1:400  
R5 - 545010 1:1000  
[30/12/2019 22:42 bilques.jacobs] Result verified

OK Cancel

Abdominal ultrasound +/- CT scan may be helpful in detecting presence of liver mass +/- intra-abdominal masses.

## Final diagnosis

?Hepatocellular carcinoma

This case allowed me to become familiar with the concepts related to limitations of an assay. Having come across the need for dilution and the concept of high-dose hook effect, I found it interesting to see the gradual increase in AFP value as further dilutions were done. These are terms and concepts that this case allowed me to become familiar with.

**Limit of Blank:** This is the highest apparent analyte concentration expected to be found when replicates of a blank sample (containing no analyte) are tested. Detects "noise" that could interfere with the result.

**Limit of Detection:** This refers to the lowest analyte

concentration likely to be reliably distinguished from the limit of blank and at which detection is feasible. LoD is determined using measured limit of blank, and test replicates known to contain a low concentration of an analyte.

**Limit of Quantitation:** This is the lowest concentration at which the analyte can not only be reliably detected but also at which some predefined goals for precision and bias are met. The LoQ may be equivalent to the LoD or it could be at a higher concentration. This is the limit that is clinically significant.

---

## 2. Creatine Kinase

Ward	Pollsmoor Female Centre	D.O.B/Age	10/10/1988
Consultant			

Creatine kinase (CK)                      265 070 H                      U/L                      20 – 180

Request form: Unable to obtain on Equation document viewer.  
Differential diagnosis in this patient includes rhabdomyolysis, severe burns, myocardial injury or ischaemia.

### History

Unavailable.

Unavailable.

Sodium		139
mmol/L	136 – 145	
Potassium		3.8
mmol/L	3.5 – 5.1	

Urea		2.6
mmol/L	2.1 – 7.1	
Creatinine		63
umol/L	49 – 90	
eGFR (MDRD formula)		>60
mL/min/1.73 m <sup>2</sup>		
Total protein		76
g/L	60 – 78	
Albumin		46
g/L	35 – 52	
Total bilirubin		6
umol/L	5 – 21	
Conjugated bilirubin (DBil)		2
umol/L	0 – 3	
Alanine transaminase (ALT)		<b>317 H</b>
U/L	7 – 35	
Aspartate transaminase (AST)		<b>1727 H</b>
U/L	13 – 35	
Alkaline phosphatase (ALP)		<b>153 H</b>
U/L	42 – 98	
Gamma-glutamyl transferase (GGT)		<b>43 H</b>
U/L	<40	

Folder unavailable. No treating doctor listed.

## Final diagnosis

Rhabdomyolysis (most likely secondary to blunt force trauma).

Creatine kinase is an enzyme primarily found in muscle tissue that catalyzes the conversion of creatine and adenosine

triphosphate (ATP) into phosphocreatine and adenosine diphosphate (ADP). This reaction is reversible and thus phosphocreatine serves as a rapidly available source of ATP. When muscle tissue is stressed or inflamed, the sarcoplasmic membrane becomes permeable and leaks cytosolic enzymes like creatine kinase into the bloodstream. The differential diagnosis of an elevated CK concentration is long and complex. Musculoskeletal trauma, myocardial injury, infections, and drug-induced myositis are the most common causes encountered in general clinical practice. Also worth noting in this patient is the AST is markedly elevated in comparison to the other liver enzymes. This, coupled with the elevated CK levels imply that quite significant muscle damage has occurred.

Making dilutions when results are appearing as having a greater than value on the analyzer is important. Creatine kinase is an enzyme that may be used to monitor the clinical condition of a patient. It may thus be useful to know exact values to be able to determine ongoing damage vs resolution. By preparing samples in dilution, this allows for a relatively accurate determination of the concentration of an analyte of interest by overcoming large reagent requirements.